



Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd



Parasitological research

doi: 10.1016/S2222-1808(16)61066-6

©2016 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

Molecular detection of *Anaplasma* species in questing ticks (ixodids) in EthiopiaSori Teshale<sup>1,2\*</sup>, Dirk Geysen<sup>2</sup>, Gobena Ameni<sup>3</sup>, Ketema Bogale<sup>4</sup>, Pierre Dorny<sup>2</sup>, Dirk Berkvens<sup>2</sup><sup>1</sup>College of Veterinary Medicine and Agriculture, Addis Ababa University, P.O. Box 34, Bishoftu, Ethiopia<sup>2</sup>Department of Biomedical Sciences, Institute of Tropical Medicine, Nationalestraat 155, Antwerp, Belgium<sup>3</sup>Akililu Lemma Institute of Pathobiology, Addis Ababa University, Addis Ababa, Ethiopia<sup>4</sup>Asela Regional Veterinary Laboratory, Asela, Ethiopia

## ARTICLE INFO

## Article history:

Received 2 Feb 2016

Received in revised form 22 Apr, 2nd

revised form 25 Apr 2016

Accepted 23 May 2016

Available online 21 Jun 2016

## Keywords:

*Anaplasma* species

Ticks

Vectors

Ethiopia

## ABSTRACT

**Objective:** To identify *Anaplasma* spp. in questing ticks with emphasis on *Anaplasma phagocytophilum* (*A. phagocytophilum*) and *Anaplasma ovis* (*A. ovis*) in Ethiopia.**Methods:** DNA extracted from 237 questing ticks [*Rhipicephalus evertsi* (*R. evertsi*) (*n* = 61), *Rhipicephalus pulchellus* (*R. pulchellus*) (*n* = 54), *Rhipicephalus decoloratus* (*n* = 1), *Amblyomma variegatum* (*n* = 22), *Amblyomma lepidum* (*n* = 36), *Amblyomma* nymphs (*n* = 6), *Amblyomma gemma* (*n* = 7) and *Hyalomma marginatum* (*Hy. marginatum*) (*n* = 53)] were tested by PCR-RFLP assay.**Results:** Overall 32 (13.33%; 95% confidence interval: 9.8%–18.3%) of the ticks were positive for *Anaplasma* spp. DNA. *Anaplasma marginale* was detected in *Hy. marginatum* and *R. pulchellus*. *Anaplasma centrale* was identified in *R. evertsi*, *R. pulchellus* and *Hy. marginatum*. *A. ovis* was detected in *R. evertsi*, *Amblyomma* spp. and *Hyalomma* spp. *A. phagocytophilum* was detected only in *R. pulchellus* and *Anaplasma* sp. *omatijenne* was detected only in *Amblyomma lepidum*. *Ehrlichia* species were not detected in any of the tick species examined.**Conclusions:** The results demonstrated the presence of several *Anaplasma* spp. including the zoonotic *A. phagocytophilum* and potentially zoonotic *A. ovis*. Our finding identified potential vectors of *A. ovis* to be further confirmed. However, an extended study is needed to identify the potential vectors of *A. phagocytophilum*. The variety of *Anaplasma* spp. identified in this study suggests risks of anaplasmosis in animals and humans in the country.

## 1. Introduction

Anaplasmosis caused by *Anaplasma marginale* (*A. marginale*) is a worldwide disease of domestic ruminants, especially cattle, and has been reported in all continents causing huge loss to cattle industry[1]. The infection of domestic ruminants with *Anaplasma centrale* (*A. centrale*) and *Anaplasma* sp. *omatijenne* (*A. sp. omatijenne*) has been also known even though these two *Anaplasma* spp. are considered non-pathogenic[1,2]. *Rhipicephalus* spp. are known to be important vectors of *A. marginale* throughout the world[1,3]. Other *Anaplasma* spp. infecting domestic ruminants such as *Anaplasma phagocytophilum* (*A. phagocytophilum*) and *Anaplasma ovis* (*A. ovis*) are either neglected or less investigated so that little is known about their epidemiology and vectors under African conditions[4,5].

According to Dulmer *et al.*[6], *A. phagocytophilum* is a recently emended species of bacteria that comprises *Ehrlichia phagocytophila*, *Ehrlichia equi* and the agent of human granulocytic anaplasmosis. It is a multihost bacterium infecting various species of wild and domestic animals and humans[7-10]. The mortality associated with *A. phagocytophilum* infection is low in animals, but significant economic losses associated with drop in milk yield, abortion and infertility and reduced weight gain have been observed in pastured animals[5,11]. Deaths have been also recorded in weaker animals if they are not treated[12]. The number of human cases associated with *A. phagocytophilum* infection has been increasing in USA, Europe, Middle East and Asia since its recognition as a human pathogen. Tick attachment, contact with infected animal blood and prenatal infection have been associated with human infections[13]. Human infections can result in severe clinical consequences with a hospitalization rate as high as 36% in USA and a mortality rate of over 26.5% in China[5].

*A. ovis* has been known to infect domestic and wild ruminants since 1912[14,15]. It is considered to be endemic in tropical and subtropical regions, but it is frequently reported in temperate

\*Corresponding author: Sori Teshale, College of Veterinary Medicine and Agriculture, Addis Ababa University, P.O. Box 34, Bishoftu, Ethiopia.

Tel: +251 911 36 36 37

E-mail: teshalesori2002@yahoo.com

The journal implements double-blind peer review practiced by specially invited international editorial board members.

regions. It has been detected in small ruminants in Europe, USA, Africa and Asia[14]. The mortality associated with the infection of *A. ovis* is not frequent, even though it causes huge financial losses to farming stock because of the reduced productivity[14,16]. It has now gained more importance as a result of observations that suggest its zoonotic importance following detection of the variant of *A. ovis* in human patients in Cyprus[17].

The vectors of *A. phagocytophilum* and *A. ovis* have been shown to vary among and within different continents and countries[5,8,17,18]. It has been also shown that *A. ovis* can be transmitted by bites of flies such as sheep ked[19]. *A. ovis* and *A. phagocytophilum* are recently identified in Ethiopia[20]. This has significant implications for Ethiopia where open range animal farming and ecotourism are the main sectors to alleviate poverty. Understanding the epidemiology of anaplasmosis caused by these *Anaplasma* spp. requires the knowledge of their tick vectors. Identification of the potential tick vectors of *A. ovis* and *A. phagocytophilum* in Ethiopia is the main objective of this study.

## 2. Materials and methods

### 2.1. Field sites for collection of ticks

Three sites where the occurrence of *Anaplasma* spp. including *A. phagocytophilum* and *A. ovis* was confirmed in domestic ruminants previously were purposely selected for collection of unfed ticks from the field for molecular analysis[20]. They were Bishoftu, Bako and Awash Depression. Bishoftu is the main town of Ada'a District in the east of Shewa Zone, Central Oromia, Ethiopia. It was located at a distance of 45 km east of Addis Ababa. Bako was a district in the west of Shewa Zone of Oromia State, Ethiopia, which was located at about 225 km away from the capital. Samples from Awash Depression were collected from three different localities (Fantale, Gari and Marti). Since these three sites were closer and had similar conditions, Fantale was taken as a representative location. The area was located in the east of Shewa Zone of Oromia State, about 190 km from the east of Addis Ababa. The Awash Depression is one of the irrigated areas in the mid-rift valley of Ethiopia. The detailed characteristics of the study sites were given in Table 1.

**Table 1**

Characteristics of the study sites where unfed ticks were collected.

Characteristics	Sites		
	Bishoftu	Bako	Awash Depression
Location	9° N, 4° E	9°8' N, 37°5' E	8°58'30" N–8°58'3" N, 39°55'48" E–39°56'00" E
Mean temperature	8.5–30.7 °C	13.5–27.9 °C	29–38 °C
Annual rainfall	1156 mm	1227 mm	560 mm
Humidity	61.30%	85.00%	Not obtained
Vegetation type	Woody vegetation	Forest type	<i>Acacia</i> woodland
Altitude	1550 m	1650 m	955–2007 m
Climate type	Intermediate	Wet, warm, humid	arid, semi-arid
Farming type	Mixed	Mixed	Livestock based
Main livestock	Cattle, sheep, goats	Cattle, sheep	Cattle, goats, camels
Production system	Commercial, smallholder	Smallholder	Smallholder

### 2.2. Collection and identification of ticks

Sampling of the ticks was carried out in September and October, 2013. The ticks were unfed, actively quested and hunted. Tick

collection was carried out by flagging vegetation on pastures and wooded areas bordering farms and homesteads as described by Uys et al.[21]. Most of the collections were carried out during morning hours. Some of the ticks were collected just while actively moving near kraals late in the afternoon when animals came back from pastures. The ticks were preserved in 70% ethanol and transported to the Veterinary Parasitology Laboratory of the College of Veterinary Medicine and Agriculture, Addis Ababa University, Bishoftu. The identification of ticks was done based on their morphological characteristics using standard identification keys described by Walker et al.[22].

### 2.3. Extraction of DNA from ticks

DNA was extracted from the ticks using the boom extraction method as described previously[23] with modifications described previously by Teshale et al.[20].

### 2.4. Amplification of DNA with PCR

A semi-nested PCR was used to amplify a fragment of about 925 bp of the 16S rDNA. Amplification was carried out using EHR 16SD (5'-GGTACCYACAGAAGAAGTCC-3')[24] and EBR3 (5'-TTGTAGTCGCCATTGTAGCAC-3')[20] primers for the first round of amplification and EHR 16SD and EBR2 (5'-TGCTGACTTGACATCATCCC-3')[20] for the second round of the reaction. The reaction mix consisted of HotStartTaq Master Mix (2.5 IU of DNA polymerase, PCR buffer containing 1.5 mmol/L MgCl<sub>2</sub> and 200 μmol/L of each deoxyribonucleotide triphosphates), 0.2 μmol/L of each primer and PCR water. The PCR reaction was carried out in a total volume of 25 μL using a programmable thermocycler (T3 thermacycler Biometra®, Westburg, NL). The PCR procedures were described by Teshale et al.[20].

All the PCR products were visualized by gel electrophoresis in Tris-acetate-EDTA buffer (0.04 mol/L Tris, 0.4 mmol/L EDTA, pH = 7.7–8.8) using 2% agarose at 100 V for 20 min and stained with ethidium bromide. Negative samples were retested at 1/10 dilution for any possible inhibition effect. Throughout the PCR procedures, PCR mix with no DNA template was used as a negative control while DNA from an *in vitro* culture of *Ehrlichia ruminantium*, *A. marginale* and *A. phagocytophilum* was used as the positive control.

### 2.5. Restriction fragment length polymorphism (RFLP) analysis of the amplified products

The amplified products from positive samples were digested by using restriction enzymes, *Mbo* II, *Hha* I and *Msp* I, to identify the species of *Anaplasma* detected as described by Teshale et al.[20]. The restriction was done in a final volume of 15 μL consisting of 4 μL DNA (PCR product) and 11 μL RFLP mix (0.3 μL per final volume of restriction enzymes, Milli-Q water and buffer for each enzyme, Biolabs, New England). Incubation was done overnight at temperatures specific for each enzyme. The restricted fragments were separated on a 2% high resolution agarose gel by electrophoresis in Tris-acetate-EDTA buffer (0.04 mol/L Tris, 0.4 mmol/L EDTA, pH = 7.7–8.8) at 100 V for 40 min and visualized under UV illumination after staining with ethidium bromide (final concentration of 0.5 μg/mL).

### 3. Results

A total of 237 ixodid ticks [*Rhipicephalus evertsi* (*R. evertsi*) ( $n = 61$ ), *Rhipicephalus pulchellus* (*R. pulchellus*) ( $n = 54$ ), *Rhipicephalus decoloratus* ( $n = 1$ ), *Amblyomma variegatum* (*A. variegatum*) ( $n = 22$ ), *Amblyomma lepidum* (*A. lepidum*) ( $n = 36$ ), *Amblyomma* nymphs ( $n = 6$ ), *Amblyomma gemma* ( $n = 7$ ) and *Hyalomma marginatum* (*Hy. marginatum*) ( $n = 53$ )] were collected from the field and analyzed for *Anaplasma* spp. DNA targeting 16S RNA gene. All ticks were tested by PCR and positive ones were subsequently analyzed by RFLP to identify the species of *Anaplasma* detected. DNA belonging to five *Anaplasma* spp. (*A. marginale*, *A. centrale*, *A. ovis*, *A. sp. omatijenne* and *A. phagocytophilum*) was identified in the ticks (Table 2).

**Table 2**

Results of molecular analysis of unfed ticks for *Anaplasma* species from selected sites in Ethiopia.

Site	Tick species	Tested number	Stage	Positive number and species
Bishoftu	<i>R. evertsi</i>	22	Adult	1 <i>A. centrale</i>
	<i>A. variegatum</i>	17	Adult	0
	<i>A. lepidum</i>	20	Adult	0
	<i>Hy. marginatum</i>	44	Adult	1 <i>A. marginale</i>
Bako	<i>R. evertsi</i>	30	Adult	6 <i>A. ovis</i>
	<i>A. lepidum</i>	9	Adult	1 <i>A. ovis</i> , 2 <i>A. sp. omatijenne</i>
	<i>Amblyomma</i>	6	Nymph	4 <i>A. ovis</i>
Awash Depression				
Fantale	<i>R. pulchellus</i>	8	Adult	1 <i>A. marginale</i> , 1 <i>A. centrale</i>
	<i>Hy. marginatum</i>	7	Adult	0
	<i>R. evertsi</i>	1	Adult	0
	<i>A. variegatum</i>	1	Adult	0
Marti	<i>R. pulchellus</i>	46	Adult	2 <i>A. phagocytophilum</i> , 4 <i>A. centrale</i> , 6 <i>A. marginale</i>
	<i>A. lepidum</i>	6	Adult	1 <i>A. ovis</i>
	<i>R. evertsi</i>	4	Adult	1 <i>A. ovis</i>
	<i>R. decoloratus</i>	1	Larva	0
	<i>A. variegatum</i>	4	Adult	0
	<i>A. gemma</i>	7	Adult	0
	Gari	<i>A. lepidum</i>	1	Adult
<i>R. evertsi</i>		1	Adult	0
<i>Hy. marginatum</i>		2	Adult	1 <i>A. ovis</i> , 1 <i>A. centrale</i>

*R. decoloratus*: *Rhipicephalus decoloratus*.

*A. marginale* was identified in 2.3% [95% confidence interval (CI): 0.1%–12%] of *Hy. marginatum* collected from Bishoftu, 12.5% (95% CI: 0.3%–2.6%) of *R. pulchellus* from Fantale and in 13% (95% CI: 4.9%–26.3%) of *R. pulchellus* from Marti. *A. centrale* was identified in 4.5% (95% CI: 0.1%–22.8%) of *R. evertsi* collected from Bishoftu, 12.5% (95% CI: 0.3%–52.6%) and 8.7% (95% CI: 2.4%–20.8%) of *R. pulchellus* from Fantale and Marti, respectively, and in 50% (95% CI: 1.3%–98.7%) of *Hy. marginatum* collected from Gari. Only two (22.2%; 95% CI: 2.8%–60.0%) of *A. lepidum* collected from Bako gave positive signals for *A. sp. omatijenne*.

The DNA of *A. phagocytophilum* was detected only in 4.3% (95% CI: 0.5%–12.7%) of *R. pulchellus* collected from Marti whereas no other ticks were found to contain DNA belonging to this pathogen. The DNA of *A. ovis* was identified in 20% (95% CI: 7.7%–38.6%) of *R. evertsi*, 11.1% (95% CI: 0.3%–48.2%) of *A. lepidum* and 66.7% (95% CI: 22.3%–95.7%) of *Amblyomma* nymphs collected from Bako. All together *A. ovis* was identified in 24.4% (95% CI: 12.9%–39.5%) of the ticks collected from Bako. Among ticks collected and tested from Marti, *A. ovis* was detected in 25% (95% CI: 0.6%–80.6%) of *R. evertsi* and 16.7% (95% CI: 0.4%–64.1%) of *A. lepidum* specimens. It was detected in 50% (95% CI: 1.3%–

98.7%) of *Hy. marginatum* collected from Gari. No ticks collected from Bishoftu contained DNA of *A. ovis*. *Ehrlichia* spp. were not identified in any of the ticks tested.

### 4. Discussion

Anaplasmosis has been well recognized in many countries of the world resulting in enormous economic losses in ruminant industry despite infrequent mortalities[1,8,17]. Variants of *Anaplasma* species have been also associated with human illness[5,17,25]. The occurrence of zoonotic *Anaplasma* spp. such as *A. ovis* and *A. phagocytophilum* was recently reported in domestic ruminants in Ethiopia[20]. However, no study has been carried out on unfed ticks in Ethiopia. To the best of our knowledge, this is the first report of the occurrence of *Anaplasma* spp. in unfed ticks collected from the field in the country. Knowledge of the ticks involved in the transmission of *Anaplasma* spp. is required for better understanding of anaplasmosis. Identification of *Anaplasma* spp. is the first step in the identification of tick vectors. In this study, we screened unfed ixodid ticks for the DNA of *Anaplasma* spp. focusing on *A. ovis* and *A. phagocytophilum* in Ethiopia.

In this study, unfed *Rhipicephalus* spp., *Hyalomma* spp. and *Amblyomma* spp. were found positive for *A. ovis* DNA. This suggests that these ticks could be involved in the transmission of *A. ovis*. The occurrence of *A. ovis* infections in resident small ruminant population was confirmed recently[20]. Now the presence of *A. ovis* in unfed tick population in the country is proved. The vector role of ticks that belong to the genus *Hyalomma* has already been documented elsewhere[26], in which unfed *Hyalomma* spp. were found to be infected with *A. ovis*. *Rhipicephalus* spp. (*R. evertsi* and *R. pulchellus*) have been also shown to be competent vectors of *Anaplasma* spp. Zhou *et al.*[1] showed that *R. evertsi* was proved to be competent vector of *A. marginale*. Therefore, the detection of DNA of *A. ovis* in unfed *Hyalomma* spp. and *Rhipicephalus* spp. in this study is in agreement with the previous reports made elsewhere in the world. In this study, *A. ovis* was also detected in two *A. lepidum* and four *Amblyomma* nymphs. *Amblyomma* spp. are mostly recognized for their role in the transmission of *Ehrlichia* spp. and *Theileria mutans*[27]. Yang *et al.*[11] have previously documented that *Amblyomma* spp. are vectors of *A. ovis* in Asia. The detection of *A. ovis* DNA in adults and nymphs that did not attach to animals suggests the possible involvement of *Amblyomma* spp. in the transmission of this bacterium. The actual role of the tick species found positive with *A. ovis* DNA in this study in the transmission of the agent to resident ruminants needs to be further confirmed by experimental studies. However, our results showed that *A. ovis* could be transmitted by several genera and species of ticks including *Amblyomma* spp. This is supported by earlier findings reported elsewhere showing that *A. ovis* is transmitted by many tick species and several biting flies[19].

Two DNA specimens from *R. pulchellus* gave positive results for *A. phagocytophilum*. This tick species was collected only from Awash Depression and its surrounding, which has typical semi-arid climate. Previous studies also showed that this tick species is found to the east of Rift Valley, which has drier weather. The occurrence of the infection with *A. phagocytophilum* was confirmed in ruminants in wetter and humid areas such as Bako where *R. pulchellus* are

not found[20]. This suggests the involvement of other tick species in the transmission of this rickettsia. An extended study covering a larger area, more tick species and quantity is needed to identify the potential vectors of *A. phagocytophilum* in the country.

In conclusion the findings of this study underline the potential for *Rhipicephalus*, *Hyalomma* and *Amblyomma* to transmit *A. ovis* among ruminants in Ethiopia. This is useful finding for small ruminant sector and could be considered during genetic improvement and translocation programs. The risks of human infections could also be highlighted as ecotourism and outdoor activities are growing in the country. Therefore, our findings could be used as preliminary information for the public health authorities. Further investigations are warranted in order to identify the vectors associated with the transmission of these *Anaplasma* spp. and elucidate the pathogenic role of *A. ovis* and *A. phagocytophilum* in human and animal health in the country.

### Conflict of interest statement

We declare that we have no conflict of interest.

### References

- [1] Zhou M, Cao SN, Sevinc F, Sevinc M, Ceylan O, Moumouni PFA, et al. Molecular detection and genetic identification of *Babesia bigemina*, *Theileria annulata*, *Theileria orientalis* and *Anaplasma marginale* in Turkey. *Ticks Tick Borne Dis* 2016; **7**(1): 126-34.
- [2] Kasari TR, Miller RS, James AM, Freier JE. Recognition of the threat of *Ehrlichia ruminantium* infection in domestic and wild ruminants in the continental United States. *J Am Vet Med Assoc* 2010; **237**(5): 520-30.
- [3] Cabezas-Cruz A, de la Fuente J. *Anaplasma marginale* major surface protein 1a: a marker of strain diversity with implications for control of bovine anaplasmosis. *Ticks Tick Borne Dis* 2015; **6**(3): 205-10.
- [4] Gharbi M, Omri H, Jedidi M, Zorii S, Darghouth MA. Epidemiological study of sheep anaplasmosis (*Anaplasma ovis* infection) in Kairouan, Central Tunisia. *J Adv Parasitol* 2015; **2**(2): 30-3.
- [5] Stuen S, Granquist EG, Silaghi C. *Anaplasma phagocytophilum*--a widespread multi-host pathogen with highly adaptive strategies. *Front Cell Infect Microbiol* 2013; **3**: 31.
- [6] Dumler JS, Barbet AF, Bekker CP, Dasch GA, Palmer GH, Ray SC, et al. Reorganisation of genera in the families of Rickettsiaceae and Anaplasmataceae in the order Rickettsiales: unification of some species of *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*, descriptions of six new species combinations and designation of *Ehrlichia equi* and 'HGE agent' as subjective synonyms of *Ehrlichia phagocytophila*. *Int J Syst Evol Microbiol* 2001; **51**(Pt 6): 2145-65.
- [7] Cabezas-Cruz A, Alberdi P, Ayllón N, Valdés JJ, Pierce R, Villar M, et al. *Anaplasma phagocytophilum* increases the levels of histone modifying enzymes to inhibit cell apoptosis and facilitate pathogen infection in the tick vector *Ixodes scapularis*. *Epigenetics* 2016; **11**(4): 303-19.
- [8] Woldehiwet Z. The natural history of *Anaplasma phagocytophilum*. *Vet Parasitol* 2010; **167**(2-4): 108-22.
- [9] Gorna M, Adaszek L, Policht K, Skrzypczak M, Winiarczyk S. Detection of *Anaplasma phagocytophilum* in a cat. *Vet Med* 2013; **58**(1): 39-43.
- [10] Ebani VV, Bertelloni F, Turchi B, Filogari D, Cerri D. Molecular survey of tick-borne pathogens in ixodid ticks collected from hunted wild animals in Tuscany, Italy. *Asian Pac J Trop Med* 2015; **8**(9): 714-7.
- [11] Yang J, Li Y, Liu Z, Liu J, Niu Q, Ren Q, et al. Molecular detection and characterization of *Anaplasma* spp. in sheep and cattle from Xinjiang, Northwest China. *Parasit Vectors* 2015; **8**: 108.
- [12] Lee SH, Jung BY, Kwak D. Evidence of *Anaplasma* spp. exposure in native Korean goats (*Capra hircus coreanae*). *Vet Med* 2015; **60**(5): 248-52.
- [13] Henningsson AJ, Wilhelmsson P, Gyllemark P, Kozak M, Matussek A, Nyman D, et al. Low risk of seroconversion or clinical disease in humans after a bite by an *Anaplasma phagocytophilum*-infected tick. *Ticks Tick Borne Dis* 2015; **6**(6): 787-92.
- [14] Mashebe P, Lyaku JR, Mause F. Occurrence of ticks and tick-borne diseases of livestock in Zambezi region: a review. *J Agric Sci* 2014; **6**(2): 142-9.
- [15] da Silva JB, André MR, Machado RZ. Low genetic diversity of *Anaplasma marginale* in calves in an endemic area for bovine anaplasmosis in the state of São Paulo, Brazil. *Ticks Tick Borne Dis* 2016; **7**(1): 20-5.
- [16] Joazeiro AC, Martins J, Masuda A, Seixas A, da Silva Vaz Junior I. A PCR for differentiate between *Anaplasma marginale* and *A. centrale*. *Acta Sci Vet* 2015; **43**: 1270.
- [17] Atif FA. *Anaplasma marginale* and *Anaplasma phagocytophilum*: Rickettsiales pathogens of veterinary and public health significance. *Parasitol Res* 2015; **114**(11): 3941-57.
- [18] Li H, Zheng YC, Ma L, Jia N, Jiang BG, Jiang RR, et al. Human infection with a novel tick-borne *Anaplasma* species in China: a surveillance study. *Lancet Infect Dis* 2015; **15**(6): 663-70.
- [19] Hornok S, de la Fuente J, Biró N, Fernández de Mera IG, Meli ML, Elek V, et al. First molecular evidence of *Anaplasma ovis* and *Rickettsia* spp. in keds (Diptera: Hippoboscidae) of sheep and wild ruminants. *Vector Borne Zoonotic Dis* 2011; **11**(10): 1319-21.
- [20] Teshale S, Geysen D, Ameni G, Asfaw Y, Berkvens D. Improved molecular detection of *Ehrlichia* and *Anaplasma* species and its application in *Amblyomma* ticks collected from cattle and sheep in Ethiopia. *Ticks Tick Borne Dis* 2015; **6**(1): 1-7.
- [21] Uys AC, Horak IG, Harrison A. Questing ixodid ticks on the vegetation of sable antelope and multi-herbivore enclosures in Thabazimbi. *J S Afr Vet Assoc* 2015; **86**: 1243.
- [22] Walker AR, Boattour A, Camicas JL, Estradar-Pena A, Latif A, Preston P. *Ticks of domestic animals in Africa, a guide to identification of species*. Edinburgh: Bioscience Reports; 2003, p. 1-227.
- [23] Herrmann BG, Frischauf AM. Isolation of genomic DNA. *Methods Enzymol* 1987; **152**: 180-3.
- [24] Hornok S, Földvári G, Elek V, Naranjo V, Farkas R, de la Fuente J. Molecular identification of *Anaplasma marginale* and rickettsial endosymbionts in blood-sucking flies (Diptera: Tabanidae, Muscidae) and hard ticks (Acari: Ixodidae). *Vet Parasitol* 2008; **154**(3-4): 354-9.
- [25] Werden L, Lindsay LR, Barker IK, Bowman J, Gonzales EK, Jardine CM. Prevalence of *Anaplasma phagocytophilum* and *Babesia microti* in *Ixodes scapularis* from a newly established lyme disease endemic area, the Thousand Islands region of Ontario, Canada. *Vector Borne Zoonotic Dis* 2015; **15**(10): 627-9.
- [26] Sarih M, M'Ghirbi Y, Bouattour A, Gern L, Baranton G, Postic D. Detection and identification of *Ehrlichia* spp. in ticks collected in Tunisia and Morocco. *J Clin Microbiol* 2005; **43**(3): 1127-32.
- [27] Estrada-Peña A, Salman M. Current limitations in the control and spread of ticks that affect livestock: a review. *Agriculture* 2013; **3**(2): 221-35.